

Total No. of Questions—8]

[Total No. of Printed Pages—6

Seat No.	
-------------	--

[5352]-511

S.E. (Mechanical/Auto/S/W) (I Sem.) EXAMINATION, 2018

ENGINEERING MATHEMATICS—III

(2015 PATTERN)

Time : Two Hours

Maximum Marks : 50

N.B. :- (i) Answer Q. No. 1 or Q. No. 2, Q. No. 3 or Q. No. 4,
Q. No. 5 or Q. No. 6, Q. No. 7 or Q. No. 8.

(ii) Neat diagrams must be drawn wherever necessary.

(iii) Figures to the right indicate full marks.

(iv) Assume suitable data, if necessary.

(v) All questions are compulsory.

1. (a) Solve any two of the following : [8]

(i) $(D^2 + 2D + 1)y = xe^{-x} \cos x$

(ii) $\frac{d^2 y}{dx^2} + 3\frac{dy}{dx} + 2y = e^{e^x}$ (using method of variation of parameter)

(iii) $(3x + 2)^2 \frac{d^2 y}{dx^2} + 3(3x + 2) \frac{dy}{dx} - 36y = 3x^2 + 4x + 1.$

(b) Using suitable Fourier transform, solve the following equation : [4]

$$\int_0^{\infty} f(x) \sin \lambda x \, dx = \begin{cases} 1 - \lambda & 0 \leq \lambda \leq 1 \\ 0 & \lambda \geq 1 \end{cases}.$$

P.T.O.

Or

2. (a) Solve any one : [4]

(i) Find Laplace transform of :

$$e^{-4t} \int_0^t \frac{\sin 3t}{t} dt.$$

(ii) Find Inverse Laplace transform of $\frac{s}{s^2 + 6s + 25}$.

(b) Using Laplace transform solve the D.E. : [4]

$$y'' + 4y' + 13y = \frac{1}{3} e^{-2t} \sin 3t, y(0) = 1, y'(0) = -2.$$

(c) A body of weight $W = 1$ N is suspended from a spring stretches it 4 cm. If the weight is pulled down 8 cm below the equilibrium position and then released : [4]

(i) Set up a differential equation.

(ii) Find the position and velocity as function of time.

3. (a) The first four moments of a distribution about the value 5 are 2, 20, 40 and 50. From the given information obtain the first four central moments and coefficient of skewness and kurtosis. [4]

(b) In a certain factory turning out razor blades, there is a small chance of $\frac{1}{500}$ for any blade to be defective. The blades are supplied in a packet of 10. Use Poisson distribution to calculate

the approximate number of packets containing : [4]

(i) no defective blades in a consignment

(ii) two defective blades

in a consignment of 10,000 packets.

(c) Find the directional derivative of the function $\phi = e^{2x - y + z}$ at (1, 1, 1) in the direction of tangent to the curve : [4]

$x = e^{-t}$, $y = 2 \sin t + 1$, $z = t - \cos t$ at $t = 0$.

Or

4. (a) Find the regression line of y on x for the following data : [4]

x	y
10	18
14	12
18	24
22	6
26	30
30	36

(b) Prove the following (any one) : [4]

(i)
$$\nabla \left[\bar{b} \cdot \nabla \left(\frac{1}{r} \right) \right] = \frac{3\bar{r}(\bar{b} \cdot \bar{r})}{r^5} - \frac{\bar{b}}{r^3}$$

(ii)
$$\nabla^2 (r^n \log r) = [n(n+1) \log r + 2n + 1] r^{n-2}$$

(c) Show that the vector field : [4]

$$\bar{F} = (2xz^3 + 6y) \bar{i} + (6x - 2yz) \bar{j} + (3x^2 z^2 - y^2) \bar{k}$$

is irrotational and hence find scalar function ϕ such that

$$\bar{F} = \nabla \phi.$$

5. (a) Find the work done in moving a particle once round the ellipse $\frac{x^2}{25} + \frac{y^2}{16} = 1$, $z = 0$ under the field of force given

by : [5]

$$\vec{F} = (2x - y + z) \vec{i} + (x + y - z^2) \vec{j} + (3x - 2y + 4z) \vec{k}.$$

- (b) Use divergence theorem to evaluate : [4]

$$\iiint_S (x \vec{i} + y \vec{j} + z^2 \vec{k}) \cdot d\vec{S}$$

where S is the curved surface of the cylinder $x^2 + y^2 = 4$, bounded by the planes $z = 0$ and $z = 2$.

- (c) Evaluate : [4]

$$\iint_S (\nabla \times \vec{F}) \cdot \hat{n} dS$$

where S is the plane surface of a lamina bounded by $x = 0$, $y = 0$, $x = 1$, $y = 1$, $z = 2$ and .

$$\vec{F} = y^2 \vec{i} + x^2 \vec{j} + z \vec{k}.$$

Or

6. (a) Evaluate : [4]

$$\int_C \vec{F} \cdot d\vec{r}$$

where

$$\vec{F} = \sin y \vec{i} + x(1 + \cos y) \vec{j}$$

and C is the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, $z = 0$.

(b) Show that : [4]

$$\iiint_V \frac{dv}{r^2} = \iint_S \frac{\bar{r} \cdot \hat{n}}{r^2} ds.$$

(c) Evaluate : [5]

$$\iint_S \text{curl } \bar{F} \cdot \hat{n} ds$$

for the surface of the paraboloid

$$z = 9 - (x^2 + y^2)$$

where

$$\bar{F} = (x^2 + y - 4) \bar{i} + 3xy \bar{j} + (2xz + z^2) \bar{k}.$$

7. (a) If $\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}$ represents the vibrations of a string of length l fixed at both ends, find the solution with boundary conditions : [7]

(i) $y(0, t) = 0$

(ii) $y(l, t) = 0$

(iii) $\left(\frac{\partial y}{\partial t}\right)_{t=0} = 0$

(iv) $y(x, 0) = kx^2, \quad 0 \leq x \leq l.$

(b) Solve $\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}$ if [6]

(i) $u(0, t) = 0$

$$(ii) \quad u_x(l, t) = 0$$

$$(iii) \quad u(x, t) \text{ is bounded}$$

$$(iv) \quad u(x, 0) = \frac{2x}{l}, \quad 0 \leq x \leq l.$$

Or

8. (a) A rectangular plate with insulated surface is 4 cm wide and so long to its width that it may be considered infinite in length.

If the temperature of the short edge $y = 0$ is given by : [6]

$$\begin{aligned} u &= 2x & 0 \leq x \leq 2 \\ &= 2(4 - x) & 2 \leq x \leq 4 \end{aligned}$$

two long edges $x = 0$, $x = 4$ as well as other short edge are kept at 0°C then find $u(x, y)$.

- (b) Use Fourier transform to solve the equation $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$

$0 < x < \infty$, $t > 0$, subject to conditions : [7]

$$(i) \quad u(0, t) = 0, \quad t > 0$$

$$(ii) \quad u(x, 0) = \begin{cases} 1 & 0 < x < 1 \\ 0 & x > 1 \end{cases}$$

$$(iii) \quad u(x, t) \text{ is bounded.}$$