

P.T.O.

2. (a) Show that the pressure at a point in a fluid at rest is the same in all directions. [6]

Or

- (b) A circular plate of 4.0 m diameter is immersed in water of density 1000 kg/m³ in such a way that the plate's greatest and least depth below free surface are 6 m and 3 m respectively. Find the total pressure on the face of the plate and the position of center of pressure.
- 3. (a) Explain the conditions of equilibrium of submerged bodies. [6]
 (b) For the flow of an incompressible fluid the velocity component in the x-direction is u = ax² + by and the velocity component if the z-direction is zero. Find the velocity component v in the y-direction such that v = 9 at y = 0. [6]

Or

1.5 m

1.0 m

4. (*a*)

A Pitot tube is inserted in flow of water having density 1000 kg/m³ as shown in figure. Assuming the coefficient of Pitot tube as 0.98; determine the following at point 1 :

water

- (1) Flow velocity.
- (2) Stagnation pressure.

[4]

[5459]-114

(b) Derive the equation :

$$\frac{dp}{\rho} + VdV + gdz = 0$$

where, p is the pressure, ρ is the density, V is the velocity of a fluid particle along a stream line, g is the acceleration due to gravity and z is difference in datum. [8]

- 5. (a) What is coefficient of velocity coefficient of contraction and coefficient of discharge for an orifice ? [6]
 - (b) An orifice of 100 mm diameter discharges water under a constant head of 4.2 m. The diameter of jet at vena contracta is 8.2 cm. If the discharge through the orifice is 40 lps, determine the hydraulic coefficients of orifice. [7]

- (1) Hydrodynamically smooth and rough boundaries.
- (2) Reynolds shear stress.
- (b) For a viscous flow through circular pipe derive the expression : [9]

$$u = (\frac{-dp}{dx}) \frac{1}{4\mu} (R^2 - r^2)$$


where, u is velocity, p is the pressure, μ is the fluid viscosity, R is the outer radius of the pipe and r is the inner radius at which the velocity distribution is obtained.

(a) What are the factors affecting the growth of boundary layer ?

7.

[4]

3



The pressure drop, Δp along a straight pipe of diameter D has been experimentally studied. It is observed for laminar flow of a given fluid and pipe, Δp varies with distance between the two points 1 and 2 as shown in figure. Assume Δp as the function of D, length L, velocity V and fluid viscosity μ . Use dimensional analysis to deduce how pressure drop, Δp varies with pipe diameter D. [9]

8. Derive an expression for displacement, momentum and energy (a)thickness with a neat sketch. [9]

Or

(b) A solid sphere of 400 mm diameter is completely immersed e on oppose oppo in a flow of sea water. The velocity of flow is 1.2 m/s and specific gravity 1.025. Calculate the drag force on the sphere assuming $C_D = 0.6$. [4]

[5459]-114

(b)

4